

CHAMAEDOREA – Ecology and Economic Botany

Nancy C. Garwood (Southern Illinois University) & Sam Bridgewater (Natural History Museum, London)

Photos by N.C. Garwood & S.G.M. Bridgewater, except where noted. Produced by N.C. Garwood, R.B. Foster, & T.S. Wachter. Support from G.&B. Moore Foundation and the UK Darwin Initiative. © Natural History Museum, London SW7 5BD UK; N.C. Garwood [ngarwood@plant.siu.edu], Dept. of Plant Biology, SIU, Carbondale, IL; S.G.M. Bridgewater [s.bridgewater@nhm.ac.uk] © Environmental & Conservation Programs, The Field Museum, Chicago, IL 60605 USA. [RRC@fmnh.org] | www.fmnh.org/plantguides/ | Rapid Color Guide #195 version 1 (2005)



1 *C. tepejilote*, the pacaya palm



2 *C. ernesti-augusti*, the 'fishtail' palm

Chamaedorea is the largest genus of palms in the Neotropics. There are 11 species in Belize. These shade-tolerant plants are often the most abundant palm in the understory of tropical forests (1, 2), where they play an important ecological role (3, 4).



3 Birds and mammals disperse the seeds



4 Thrips, beetles & wind pollinate the flowers
photo: H. Porter-Morgan (NYBG)

Pacaya, the immature male inflorescences of *C. tepejilote*, has been eaten for millennia in the Mayan region (5).

Cultivars have been selected by local people as a perennial vegetable crop. It is now exported abroad (below).



5 Pacaya for sale in the Cayo market



6 *C. seifrizii* at a hotel in Cayo District



7 *C. seifrizii* being sold in Belmopan

Chamaedorea palms are used locally and internationally as indoor potted plants and in landscaping (6, 7). Only a few species used in the international horticultural trade produce seeds outside of their native range, perhaps because pollinators are not available or environmental conditions are inappropriate.



8 Cut leaves in bouquet in New York City
photo: H. Porter-Morgan (NYBG)

The cut leaves of several species are also sold for use in floral displays (8). The combined international horticultural and floral trade in *Chamaedorea* palms is a large and important industry. Seeds to support the horticultural industry (9), and cut leaves for the floral trade (10), are harvested from plants growing wild in tropical forests or in plantations and then exported.



9 Seed for export from a Belizean plantation

In Belize and Guatemala, the cut leaves and the three palm species providing them are called xaté (pronounced sha-tay). Large amounts of xaté are now being illegally and unsustainably harvested from the forests of Belize (11). Unsustainable harvesting (12) can kill individual palms and cause regional extinction of species.



10 Cut leaf bundles of xaté macho, *C. oblongata*

This activity deprives Belize of economic benefits from its natural resources, now and in the future. Other activities of the illegal collectors (called xateros), such as hunting, threatens endangered wildlife and degrades the forest ecosystem. This in turn negatively impacts other economic activities such as eco-tourism.



11 Leaf bundles of illegally cut 'fishtail' palm



12 'Fishtail' palm with all of its leaves cut off



13 Nursery of 'fishtail' seedlings for plantations

If wild harvesting of xaté in Belize proves too difficult to regulate, plantation-grown xaté may be a viable alternative (13). Several NGOs are now working with local communities to determine whether cultivation of xaté can help small farmers economically by diversifying their income base.

Palms of Belize

CHAMAEDOREA – Species

Nancy C. Garwood (Southern Illinois University) & **Sam Bridgewater** (Natural History Museum, London)

Photos by N.C. Garwood & S.G.M. Bridgewater, except where noted. Produced by N.C. Garwood, R.B. Foster, & T.S. Wachter. Support from G.&B. Moore Foundation and the UK Darwin Initiative.
© Natural History Museum, London SW7 5BD UK; N.C. Garwood [ngarwood@plant.siu.edu], Dept. of Plant Biology, SIU, Carbondale, IL; S.G.M. Bridgewater [s.bridgewater@nhm.ac.uk]
© Environmental & Conservation Programs, The Field Museum, Chicago, IL 60605 USA. [RRC@fmnh.org] | www.fmnh.org/plantguides/ | Rapid Color Guide #195 version 1 (2005)

C. ernesti-augusti (fishtail, xaté)

Height: to 2 m
Stem width: to 1.5 cm
Habit: solitary
Habitat: forest, often on limestone
Distribution: Belize, Cayo, Stan Creek, Toledo
Uses: leaves cut for floral trade, ornamental
(14-17, 2, 11-13)



14 Habit



15 Male inflorescence



16 Flowers: female (above) male (below)



17 Inflorescence with ripe fruit

C. elegans (parlor palm, xaté hembra)

Height: to 2 m
Stem width: 1.5 cm
Habit: solitary
Habitat: forest, often on limestone
Distribution: Cayo
Uses: leaves cut for floral trade, horticulture
(18-21, 8-9)



18 Habit



19 Leaf



20 Male inflorescence and flowers



21 Inflorescence with ripe fruit

C. oblongata (xaté macho, jade)

Height: to 3 m
Stem width: to 2.5 cm
Habit: solitary
Habitat: lowland forest
Distribution: Belize, Cayo, Orange Walk, Stann Creek, Toledo
Uses: leaves cut for floral trade, ornamental
ID: similar to *C. neurochlamys* (see below)
(22-25, 3, 10)



22 Habit



23 Green apex of sheaths (above); leaflets (below)



24 Flowers: male (right) female (left)



25 Inflorescence with green, unripe fruit

C. neurochlamys (false jade, monkey-tail)

Height: to 4.5 m
Stem width: to 2.5 cm
Habit: solitary
Habitat: lowland forest
Distribution: Cayo, Orange Walk, Stan Creek, Toledo
Uses: ornamental
ID: similar to *C. oblongata*, but with white apex of leaf sheath (not green) and orange unripe fruit (not green)
(26-29)



26 Habit



27 White apex of sheaths (above); leaflets (below)



28 Male inflorescence and male flowers



29 Inflorescence with ripe and orange unripe fruit

Palms of Belize

CHAMAEDOREA – Species

Nancy C. Garwood (Southern Illinois University) & Sam Bridgewater (Natural History Museum, London)

Photos by N.C. Garwood & S.G.M. Bridgewater, except where noted. Produced by N.C. Garwood, R.B. Foster, & T.S. Wachter. Support from G.&B. Moore Foundation and the UK Darwin Initiative.
© Natural History Museum, London SW7 5BD UK; N.C. Garwood [ngarwood@plant.siu.edu], Dept. of Plant Biology, SIU, Carbondale, IL; S.G.M. Bridgewater [s.bridgewater@nhm.ac.uk]
© Environmental & Conservation Programs, The Field Museum, Chicago, IL 60605 USA. [RRC@fmnh.org] | www.fmnh.org/plantguides/ | Rapid Color Guide #195 version 1 (2005)

C. tepejilote (pacaya)

Height: to 7 m
Stem width: to 10 cm
Habit: usually clonal, clumps loose or dense
Habitat: forest, often on limestone
Distribution: Cayo, Orange Walk, Stann Creek, Toledo
Uses: horticulture; male inflorescences cooked
(30-33, 1, 4-5)



30 Habit



31 Clumped stems (above) and leaflets (below)



32 Male inflorescences



33 Inflorescence with maturing fruit

C. seifrizii (bamboo palm)

Height: to 3 m
Stem width: to 2 cm
Habit: clonal, clumps dense
Habitat: open woodland or forest, often on limestone
Distribution: Belize, Cayo, Corozal, Orange Walk
Uses: horticulture
ID: differs from *C. schippii* in tight clumping of stems and lowland habitat
(34-37, 6-7)



34 Habit



35 Leaves



36 Male inflorescence and flowers



37 Inflorescence with ripe fruits

C. schippii (chapai)

Height: to 4 m
Stem width: to 3 cm
Habit: clonal, clumps loose
Habitat: forest, summits of limestone hills
Distribution: Cayo, Toledo
Uses: leaves medicinal
ID: differs from *C. seifrizii* in loose clumping of stems and hilltop habitat
(38-41)



38 Habit



39 Leaves



40 Male inflorescences and male flowers



41 Inflorescence with ripe fruits

C. adscendens

Height: to 2.5 m
Stem width: to 1 cm
Habit: solitary
Habitat: forest, summits of steep limestone hills
Distribution: Toledo
Uses: ornamental
(42-45)



42 Habit



43 Adult (above) and juvenile (below) leaves



44 Female inflorescence



45 Inflorescence with maturing fruit

Palms of Belize

CHAMAEDOREA – Species

Nancy C. Garwood (Southern Illinois University) & Sam Bridgewater (Natural History Museum, London)

Photos by N.C. Garwood & S.G.M. Bridgewater, except where noted. Produced by N.C. Garwood, R.B. Foster, & T.S. Wachter. Support from G.&B. Moore Foundation and the UK Darwin Initiative.
 © Natural History Museum, London SW7 5BD UK; N.C. Garwood [ngarwood@plant.siu.edu], Dept. of Plant Biology, SIU, Carbondale, IL; S.G.M. Bridgewater [s.bridgewater@nhm.ac.uk]
 © Environmental & Conservation Programs, The Field Museum, Chicago, IL 60605 USA. [RRC@fmnh.org] | www.fmnh.org/plantguides/ | Rapid Color Guide #195 version 1 (2005)

C. geonomiformis

Height: to 2 m
 Stem width: to 1 cm
 Habit: solitary
 Habitat: wet forest, often on limestone
 Distribution: Toledo
 Uses: ornamental
 ID: Leaves narrower & less deeply notched than *C. ernesti-augusti* (46-49)



46 Habit



47 Leaves



48 Male Inflorescence



49 Infructescence

C. arenbergiana

Height: to 4 m
 Stem width: to 3 cm
 Habit: solitary
 Habitat: wet forest
 Distribution: Toledo
 Uses: ornamental
 ID: Differing from *C. oblongata* & *C. neurochlamys* by much larger leaflets, unbranched female inflorescences, and densely packed flowers and fruits (50-52)



50 Habit



51 Adult leaf



52 Infructescence

This Guide has been produced with assistance from the UK Darwin Initiative



Glossary:

Dioecious – male & female flowers on separate plants
 Monoecious – male & female flowers on same plant
 Pinnate leaf – leaves compound, with leaflets arranged like a feather
 Bifid leaf – leaves simple, apex notched
 Petiole – stalk of leaf below leaf blade
 Rachis – central stalk of pinnate leaf with leaflets
 Inflorescence: stalk bearing flowers
 Infructescence: stalk bearing fruits

References:

Balick, MJ., MH Nee, & DE Atha. 2001. *Checklist of the Vascular Plants of Belize*.
 Henderson, A, G Galeano, & R Bernal. 1995. *Field Guide to the Palms of the Americas*. Princeton University Press.
 Hodel, D. 1992. *Chamaedorea Palms: The species and their cultivation*. International Palm Society, Lawrence, Kansas.

Taxonomic notes:

1) *C. neurochlamys* is sometimes included within the widespread *C. pinnatifrons*
 2) *C. schippii* is now considered distinct from the Costa Rican *C. graminifolia*
 3) *C. woodsoniana* (right, 53): one collection reported from northern Cayo, but identification questioned by some researchers; similar to *C. tepejilote* in size, but with narrower leaflets, solitary habit and orange unripe fruit (not green)



53 *C. woodsoniana* Habit

photo: R. Foster (F)

Key morphological characters of *Chamaedorea* in Belize

Small understory palms (1 to 12 m tall)
 Spines absent
 Dioecious
 Stems and leaf sheaths green with contrasting pale leaf scars circling stem
 Leaves bifid or pinnate, usually smooth (not pubescent or scaly)
 Leaf margins usually toothed
 Petioles often with yellow line on underside
 Flowers solitary or in groups of 2-3
 Seed surface not brain-like (usually fibrous)

Key characters of other small spineless understory palms in Belize

All are monoecious (and have male and female flowers on the same inflorescence)

If stems and leaf sheaths green and adult leaves pinnate and smooth:

Synecanthus: seed surface smooth & brain-like; inflorescence large and loosely broom-like; petiole lacking yellow line below; flowers in 6-14 lines on inflorescence branches

If stems and/or leaf sheaths brown or grayish and adult leaves bifid or pinnate:

Reinhardtia: leaves pinnate with ragged apex and 'windows' in leaf blade along rachis; flowers on surface of inflorescence (not in pits)

Geonomid palms: flowers in small pits on inflorescence; leaves without 'windows'

Aenoma: leaves bifid or pinnate; brown scales or hairs on leaves or inflorescences

Asterogyne: leaves bifid; with skirt of persistent dead leaves below crown

Calyptogyne: leaves unequally pinnate & inflorescence spikes held well above leaves